

## **Botox Worksheet**

ue	:nt:	
]	Botox <b>E</b>	Date:
1.	Medica	ation Review
	a.	Blood thinners- Eliquis, Coumadin/ Warfarin, Plavix, Aspirin, NSAIDs
		MUST BE STOPPED 3 DAYS PRIOR. You will need clearance from prescribing
		MD to stop the med prior to EACH BOTOX procedure.
		Clearance needed? YES NO from Doctor
	b.	STOP bladder meds (oxybutynin, Myrbetriq) the day before procedure. Do not
		resume it. We want to see the effects of the Botox during this time.
	c.	Antibiotic will be started 3 days BEFORE procedure. Take 1 on day OF procedure
		and continue for 3 days AFTER (total of 7 days). (Bactrim, Nitrofurantoin, Cipro)
	d.	Review Allergies- Lidocaine, Iodine (topical), antibiotics
		Allergies:

## 2. Procedure Day

- a. Upon arrival, you will empty your bladder at the office.
- b. A catheter will be inserted into your bladder. Lidocaine will be infused and allowed to sit for 30 minutes.
- c. Your provider will then cleanse your perineum with iodine and insert a camera into your bladder to view the inside of the bladder for any abnormalities.
- d. 20 small injections will be given inside the bladder prior to removal of camera.

## 3. Post Procedure Expectations

- a. It is normal for your urine to be blood tinged and have burning with urination.
- b. You may resume your blood thinners today (if applicable).
- c. Leakage may still occur and may worsen. The effects of the injections will reach its maximum effects by 3 weeks.
- d. Complete a 1-day diary **BEFORE** your 3-week follow-up visit. Keep your speci-hat for future use (will need to do diary before next Botox injection as well).
- e. At your follow-up visit, we will have you urinate in the office before your exam. We will scan your bladder to confirm your post-void residual (PVR).
- f. A minimum of 12 weeks is needed between injections. On average, Botox injections are repeated every 6-12 months, but is unique to each patient.